


WHAT IS CLAIMED IS:

1. A compound of Formula I:



- 5 or a pharmaceutically acceptable salt or stereoisomer thereof,

wherein:

- a is 0 or 1;
 10 b is 0 or 1;
 m is 0, 1, or 2;
 n is 0, 1, 2 or 3;
 r is 0 or 1;
 s is 0 or 1;
 15 t is 0, 1 or 2;
 u is 0, 1, or 2;

R¹ and R² are independently selected from: H, (C₁-C₆)alkyl, aryl, heterocyclyl and (C₃-C₆)cycloalkyl, optionally substituted with one, two or three substituents selected from R⁷;

20

R³ is selected from:

- 1) Hydrogen,
 - 2) C₁-C₁₀ alkyl;
 - 3) C₁-C₁₀ alkyl-O-R^d,
 - 4) C₂-C₁₀ alkenyl-O-R^d,
 - 5) C₂-C₁₀ alkynyl-O-R^d,
 - 6) (C₁-C₆-alkylene)_nC₃-C₈ cycloalkyl-O-R^d,
 - 7) C₁-C₁₀ alkyl-(C=O)_b-NR^cR^{c'},
 - 8) C₂-C₁₀ alkenyl-(C=O)_b-NR^cR^{c'},
 - 9) C₂-C₁₀ alkynyl-(C=O)_b-NR^cR^{c'},
 - 10) (C₁-C₆-alkylene)_nC₃-C₈ cycloalkyl-(C=O)_b-NR^cR^{c'},
 - 11) C₁-C₁₀ alkyl-S(O)_m-R^d,
 - 12) C₂-C₁₀ alkenyl-S(O)_m-R^d,
 - 13) C₂-C₁₀ alkynyl-S(O)_m-R^d,
 - 14) (C₁-C₆-alkylene)_nC₃-C₈ cycloalkyl-S(O)_m-R^d,
- 15 said alkyl, alkenyl, alkynyl and cycloalkyl are optionally substituted with one or more substituents selected from R⁶;

R⁴ is independently selected from:

- 1) (C=O)_aO_bC₁-C₁₀ alkyl,
- 2) (C=O)_aO_baryl,
- 3) CO₂H,
- 4) halo,
- 5) CN,
- 6) OH,
- 7) O_bC₁-C₆ perfluoroalkyl,
- 8) O_a(C=O)_bNR⁸R⁹,
- 9) S(O)_mR^a,
- 10) S(O)₂NR⁸R⁹,
- 11) -OPO(OH)₂;

30 said alkyl, aryl, alkenyl, alkynyl, heterocyclyl, and cycloalkyl optionally substituted with one, two or three substituents selected from R⁷;

R⁵ is selected from:

- 1) hydrogen;
- 2) (C=O)_aO_bC₁-C₁₀ alkyl,

- 5
- 3) $(\text{C}=\text{O})_a\text{O}_b\text{aryl}$,
 - 4) CO_2H ,
 - 5) halo,
 - 6) CN ,
 - 7) OH ,
 - 8) $\text{O}_b\text{C}_1\text{-C}_6$ perfluoroalkyl,
 - 9) $\text{O}_a(\text{C}=\text{O})_b\text{NR}^8\text{R}^9$,
 - 10) $\text{S}(\text{O})_m\text{R}^a$,
 - 11) $\text{S}(\text{O})_2\text{NR}^8\text{R}^9$,
 - 10 12) $-\text{OPO}(\text{OH})_2$;

said alkyl, aryl, alkenyl, alkynyl, heterocyclyl, and cycloalkyl optionally substituted with one, two or three substituents selected from R^7 ;

R^6 is independently selected from:

- 15
- 1) $(\text{C}=\text{O})_a\text{O}_b\text{C}_1\text{-C}_{10}$ alkyl,
 - 2) $(\text{C}=\text{O})_a\text{O}_b\text{aryl}$,
 - 3) $\text{C}_2\text{-C}_{10}$ alkenyl,
 - 4) $\text{C}_2\text{-C}_{10}$ alkynyl,
 - 5) $(\text{C}=\text{O})_a\text{O}_b$ heterocyclyl,
 - 20 6) CO_2H ,
 - 7) halo,
 - 8) CN ,
 - 9) OH ,
 - 10) $\text{O}_b\text{C}_1\text{-C}_6$ perfluoroalkyl,
 - 25 11) $\text{O}_a(\text{C}=\text{O})_b\text{NR}^8\text{R}^9$,
 - 12) $\text{S}(\text{O})_m\text{R}^a$,
 - 13) $\text{S}(\text{O})_2\text{NR}^8\text{R}^9$,
 - 14) oxo,
 - 15) CHO ,
 - 30 16) $(\text{N}=\text{O})\text{R}^8\text{R}^9$, or
 - 17) $(\text{C}=\text{O})_a\text{O}_b\text{C}_3\text{-C}_8$ cycloalkyl,
 - 18) $-\text{OPO}(\text{OH})_2$;

said alkyl, aryl, alkenyl, alkynyl, heterocyclyl, and cycloalkyl optionally substituted with one, two or three substituents selected from R^7 ;

R⁷ is selected from:

- 1) (C=O)_rO_s(C₁-C₁₀)alkyl,
- 2) O_r(C₁-C₃)perfluoroalkyl,
- 3) oxo,
- 5 4) OH,
- 5) halo,
- 6) CN,
- 7) (C₂-C₁₀)alkenyl,
- 8) (C₂-C₁₀)alkynyl,
- 10 9) (C=O)_rO_s(C₃-C₆)cycloalkyl,
- 10) (C=O)_rO_s(C₀-C₆)alkylene-aryl,
- 11) (C=O)_rO_s(C₀-C₆)alkylene-heterocyclyl,
- 12) (C=O)_rO_s(C₀-C₆)alkylene-N(R^b)₂,
- 13) C(O)R^a,
- 15 14) (C₀-C₆)alkylene-CO₂R^a,
- 15) C(O)H,
- 16) (C₀-C₆)alkylene-CO₂H,
- 17) (C=O)_rN(R^b)₂,
- 18) S(O)_mR^a,
- 20 19) S(O)₂N(R^b)₂, and
- 20) -OPO(OH)₂;

said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, alkylene and heterocyclyl is optionally substituted with up to three substituents selected from R^b, OH, (C₁-C₆)alkoxy, halogen, CO₂H, CN, O(C=O)C₁-C₆ alkyl, oxo, NO₂ and N(R^b)₂;

R⁸ and R⁹ are independently selected from:

- 1) H,
- 2) (C=O)O_bC₁-C₁₀ alkyl,
- 3) (C=O)O_bC₃-C₈ cycloalkyl,
- 30 4) (C=O)O_baryl,
- 5) (C=O)O_bheterocyclyl,
- 6) C₁-C₁₀ alkyl,
- 7) aryl,
- 8) C₂-C₁₀ alkenyl,
- 35 9) C₂-C₁₀ alkynyl,

- 10) heterocyclyl,
- 11) C₃-C₈ cycloalkyl,
- 12) SO₂R^a, and
- 13) (C=O)NR^b₂,

5 said alkyl, cycloalkyl, aryl, heterocyclyl, alkenyl, and alkynyl is optionally substituted with one, two or three substituents selected from R⁷, or

R⁸ and R⁹ can be taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 3-7 members in each ring and optionally containing, in
 10 addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with one, two or three substituents selected from R⁷;

R¹⁰ is selected from: H and F;

15 R¹¹ and R¹² are independently selected from: F and -CH₂F;

R¹³ and R¹⁴ are independently selected from: H and -CH₂F;

20 R^{ox} is absent or is oxo;

R^a is independently selected from: (C₁-C₆)alkyl, (C₃-C₆)cycloalkyl, aryl, or heterocyclyl, optionally substituted with one, two or three substituents selected from R⁷;

25 R^b is independently selected from: H, (C₁-C₆)alkyl, aryl, heterocyclyl, (C₃-C₆)cycloalkyl, (C=O)OC₁-C₆ alkyl, (C=O)C₁-C₆ alkyl, (C=O)aryl, (C=O)heterocyclyl, (C=O)NR^eRe' or S(O)₂R^a, optionally substituted with one, two or three substituents selected from R⁷;

30 R^c and R^{c'} are independently selected from: H, (C₁-C₆)alkyl, aryl, NH₂, OH, OR^a, -(C₁-C₆)alkyl-OH, -(C₁-C₆)alkyl-O-(C₁-C₆)alkyl, (C=O)OC₁-C₆ alkyl, (C=O)C₁-C₆ alkyl, (C=O)aryl, (C=O)heterocyclyl, (C=O)NR^eRe', S(O)₂R^a and -(C₁-C₆)alkyl-N(R^b)₂, wherein the alkyl is optionally substituted with one, two or three substituents selected from R⁷; or

35 R^c and R^{c'} can be taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 3-7 members in each ring and optionally containing, in


addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with one, two or three substituents selected from R⁷;

- 5 R^d is selected from: H, (C₁-C₆)alkyl, -(C₂-C₆)alkyl-OH, -(C₁-C₆)alkyl-O-(C₁-C₆)alkyl and -(C₁-C₆)alkyl-N(R^b)₂, wherein the alkyl is optionally substituted with one, two or three substituents selected from R⁷;

- 10 R^e and R^{e'} are independently selected from: H, (C₁-C₆)alkyl, aryl, heterocyclyl and (C₃-C₆)cycloalkyl, optionally substituted with one, two or three substituents selected from R⁷; or

- 15 R^e and R^{e'} can be taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 3-7 members in each ring and optionally containing, in addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with one, two or three substituents selected from R⁷

2. The compound according to Claim 1 of Formula II:



or a pharmaceutically acceptable salt or stereoisomer thereof,

wherein:

- a is 0 or 1;
 b is 0 or 1;
 5 m is 0, 1, or 2;
 n is 0, 1, 2 or 3;
 r is 0 or 1;
 s is 0 or 1;
 t is 0 or 1;
 10 u is 0 or 1;

R¹ and R² are independently selected from: H, (C₁-C₆)alkyl, aryl, heterocyclyl and (C₃-C₆)cycloalkyl, optionally substituted with one, two or three substituents selected from R⁷;

15 R³ is selected from:

- 1) hydrogen;
- 2) C₁-C₁₀ alkyl;
- 3) C₁-C₁₀ alkyl-O-R^d,
- 4) C₂-C₁₀ alkenyl-O-R^d,
- 20 5) C₂-C₁₀ alkynyl-O-R^d,
- 6) (C₁-C₆-alkylene)_nC₃-C₈ cycloalkyl-O-R^d,
- 7) C₁-C₁₀ alkyl-(C=O)_b-NR^cR^{c'},
- 8) C₂-C₁₀ alkenyl-(C=O)_bNR^cR^{c'},
- 9) C₂-C₁₀ alkynyl-(C=O)_bNR^cR^{c'},
- 25 10) (C₁-C₆-alkylene)_nC₃-C₈ cycloalkyl-(C=O)_bNR^cR^{c'},
- 11) C₁-C₁₀ alkyl-S(O)_m-R^d,
- 12) C₂-C₁₀ alkenyl-S(O)_m-R^d,
- 13) C₂-C₁₀ alkynyl-S(O)_m-R^d,
- 14) (C₁-C₆-alkylene)_nC₃-C₈ cycloalkyl-S(O)_m-R^d,

30 said alkyl, alkenyl, alkynyl and cycloalkyl are optionally substituted with one or more substituents selected from R⁶;

R⁴ is independently selected from:

- 35 1) (C=O)_aO_bC₁-C₁₀ alkyl,

- 2) $(\text{C}=\text{O})_a\text{O}_b\text{aryl}$,
 3) CO_2H ,
 4) halo,
 5) CN ,
 5 6) OH ,
 7) $\text{O}_b\text{C}_1\text{-C}_6$ perfluoroalkyl,
 8) $\text{O}_a(\text{C}=\text{O})_b\text{NR}^8\text{R}^9$,
 9) $\text{S}(\text{O})_m\text{R}^a$,
 10) $\text{S}(\text{O})_2\text{NR}^8\text{R}^9$, and
 10 11) $-\text{OPO}(\text{OH})_2$;

said alkyl, aryl, alkenyl, alkynyl, heterocyclyl, and cycloalkyl optionally substituted with one, two or three substituents selected from R^7 ;

R^5 is selected from:

- 15 1) hydrogen;
 2) $(\text{C}=\text{O})_a\text{O}_b\text{C}_1\text{-C}_{10}$ alkyl,
 3) $(\text{C}=\text{O})_a\text{O}_b\text{aryl}$,
 4) CO_2H ,
 5) halo,
 20 6) CN ,
 7) OH ,
 8) $\text{O}_b\text{C}_1\text{-C}_6$ perfluoroalkyl,
 9) $\text{O}_a(\text{C}=\text{O})_b\text{NR}^8\text{R}^9$,
 10) $\text{S}(\text{O})_m\text{R}^a$,
 25 11) $\text{S}(\text{O})_2\text{NR}^8\text{R}^9$,

said alkyl, aryl, alkenyl, alkynyl, heterocyclyl, and cycloalkyl optionally substituted with one, two or three substituents selected from R^7 ;

R^6 is independently selected from:

- 30 1) $(\text{C}=\text{O})_a\text{O}_b\text{C}_1\text{-C}_{10}$ alkyl,
 2) $(\text{C}=\text{O})_a\text{O}_b\text{aryl}$,
 3) $\text{C}_2\text{-C}_{10}$ alkenyl,
 4) $\text{C}_2\text{-C}_{10}$ alkynyl,
 5) $(\text{C}=\text{O})_a\text{O}_b$ heterocyclyl,
 35 6) CO_2H ,

- 7) halo,
- 8) CN,
- 9) OH,
- 10) $O_bC_1-C_6$ perfluoroalkyl,
- 5 11) $O_a(C=O)_bNR^8R^9$,
- 12) $S(O)_mR^a$,
- 13) $S(O)_2NR^8R^9$,
- 14) oxo,
- 15) CHO,
- 10 16) $(N=O)R^8R^9$, or
- 17) $(C=O)_aO_bC_3-C_8$ cycloalkyl, and
- 18) $-OPO(OH)_2$;

said alkyl, aryl, alkenyl, alkynyl, heterocyclyl, and cycloalkyl optionally substituted with one, two or three substituents selected from R^7 ;

15

R^7 is selected from:

- 1) $(C=O)_rO_s(C_1-C_{10})$ alkyl,
- 2) $O_r(C_1-C_3)$ perfluoroalkyl,
- 3) oxo,
- 20 4) OH,
- 5) halo,
- 6) CN,
- 7) (C_2-C_{10}) alkenyl,
- 8) (C_2-C_{10}) alkynyl,
- 25 9) $(C=O)_rO_s(C_3-C_6)$ cycloalkyl,
- 10) $(C=O)_rO_s(C_0-C_6)$ alkylene-aryl,
- 11) $(C=O)_rO_s(C_0-C_6)$ alkylene-heterocyclyl,
- 12) $(C=O)_rO_s(C_0-C_6)$ alkylene- $N(R^b)_2$,
- 13) $C(O)R^a$,
- 30 14) (C_0-C_6) alkylene- CO_2R^a ,
- 15) $C(O)H$,
- 16) (C_0-C_6) alkylene- CO_2H ,
- 17) $C(O)N(R^b)_2$,
- 18) $S(O)_mR^a$,
- 35 19) $S(O)_2N(R^b)_2$; and

20) $-\text{OPO}(\text{OH})_2$;

said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, alkylene and heterocyclyl is optionally substituted with up to three substituents selected from R^b , OH, $(\text{C}_1\text{-C}_6)\text{alkoxy}$, halogen, CO_2H , CN, $\text{O}(\text{C}=\text{O})\text{C}_1\text{-C}_6$ alkyl, oxo, NO_2 and $\text{N}(\text{R}^b)_2$;

5

R^8 and R^9 are independently selected from:

- 1) H,
- 2) $(\text{C}=\text{O})\text{O}_b\text{C}_1\text{-C}_{10}$ alkyl,
- 3) $(\text{C}=\text{O})\text{O}_b\text{C}_3\text{-C}_8$ cycloalkyl,
- 10 4) $(\text{C}=\text{O})\text{O}_b\text{aryl}$,
- 5) $(\text{C}=\text{O})\text{O}_b\text{heterocyclyl}$,
- 6) $\text{C}_1\text{-C}_{10}$ alkyl,
- 7) aryl,
- 8) $\text{C}_2\text{-C}_{10}$ alkenyl,
- 15 9) $\text{C}_2\text{-C}_{10}$ alkynyl,
- 10) heterocyclyl,
- 11) $\text{C}_3\text{-C}_8$ cycloalkyl,
- 12) SO_2R^a , and
- 13) $(\text{C}=\text{O})\text{NR}^b_2$,

20 said alkyl, cycloalkyl, aryl, heterocyclyl, alkenyl, and alkynyl is optionally substituted with one, two or three substituents selected from R^7 , or

R^8 and R^9 can be taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 3-7 members in each ring and optionally containing, in
 25 addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with one, two or three substituents selected from R^7 ;

R^{11} and R^{12} are independently selected from: F and $-\text{CH}_2\text{F}$;

30

R^{13} and R^{14} are independently selected from: H and $-\text{CH}_2\text{F}$, provided that when t is 1, R^{14} is H; and when u is 1, R^{13} is H;

R^{ox} is absent or is oxo;

35

R^a is independently selected from: (C₁-C₆)alkyl, (C₃-C₆)cycloalkyl, aryl, or heterocyclyl, optionally substituted with one, two or three substituents selected from R⁷;

5 R^b is independently selected from: H, (C₁-C₆)alkyl, aryl, heterocyclyl, (C₃-C₆)cycloalkyl, (C=O)OC₁-C₆ alkyl, (C=O)C₁-C₆ alkyl, (C=O)aryl, (C=O)heterocyclyl, (C=O)NR^eR^{e'} or S(O)₂R^a, optionally substituted with one, two or three substituents selected from R⁷;

10 R^c and R^{c'} are independently selected from: H, (C₁-C₆)alkyl, aryl, NH₂, OH, OR^a, -(C₁-C₆)alkyl-OH, -(C₁-C₆)alkyl-O-(C₁-C₆)alkyl, (C=O)OC₁-C₆ alkyl, (C=O)C₁-C₆ alkyl, (C=O)aryl, (C=O)heterocyclyl, (C=O)NR^eR^{e'}, S(O)₂R^a and -(C₁-C₆)alkyl-N(R^b)₂, wherein the alkyl is optionally substituted with one, two or three substituents selected from R⁷; or


15 R^c and R^{c'} can be taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 3-7 members in each ring and optionally containing, in addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with one, two or three substituents selected from R⁷;

20 R^d is selected from: H, (C₁-C₆)alkyl, -(C₂-C₆)alkyl-OH, -(C₁-C₆)alkyl-O-(C₁-C₆)alkyl and -(C₁-C₆)alkyl-N(R^b)₂, wherein the alkyl is optionally substituted with one, two or three substituents selected from R⁷;

25 R^e and R^{e'} are independently selected from: H, (C₁-C₆)alkyl, aryl, heterocyclyl and (C₃-C₆)cycloalkyl, optionally substituted with one, two or three substituents selected from R⁷; or

30 R^e and R^{e'} can be taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 3-7 members in each ring and optionally containing, in addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with one, two or three substituents selected from R⁷.

3. The compound according to Claim 2 of the Formula III:



III

or a pharmaceutically acceptable salt or stereoisomer thereof,

wherein:

5

- a is 0 or 1;
- b is 0 or 1;
- m is 0, 1, or 2;
- n is 0, 1 or 2;
- 10 r is 0 or 1;
- s is 0 or 1;
- t is 0 or 1;

R^1 and R^2 are independently selected from: H, (C_1-C_6) alkyl, aryl and (C_3-C_6) cycloalkyl,
 15 optionally substituted with one, two or three substituents selected from R^7 ;

R^4 is independently selected from:

- 4) halo,
- 5) OH,
- 20 6) $O_bC_1-C_6$ perfluoroalkyl,

R^5 is selected from:

- 1) hydrogen,
- 2) halo,
- 3) OH,
- 4) $O_bC_1-C_6$ perfluoroalkyl,

5

R^7 is selected from:

- 1) $(C=O)_rO_s(C_1-C_{10})$ alkyl,
- 2) $O_r(C_1-C_3)$ perfluoroalkyl,
- 3) oxo,
- 10 4) OH,
- 5) halo,
- 6) CN,
- 7) (C_2-C_{10}) alkenyl,
- 8) (C_2-C_{10}) alkynyl,
- 15 9) $(C=O)_rO_s(C_3-C_6)$ cycloalkyl,
- 10) $(C=O)_rO_s(C_0-C_6)$ alkylene-aryl,
- 11) $(C=O)_rO_s(C_0-C_6)$ alkylene-heterocyclyl,
- 12) $(C=O)_rO_s(C_0-C_6)$ alkylene- $N(R^b)_2$,
- 13) $C(O)R^a$,
- 20 14) (C_0-C_6) alkylene- CO_2R^a ,
- 15) $C(O)H$,
- 16) (C_0-C_6) alkylene- CO_2H , and
- 17) $C(O)N(R^b)_2$,
- 18) $S(O)_mR^a$, and
- 25 19) $S(O)_2N(R^b)_2$;

said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, alkylene and heterocyclyl is optionally substituted with up to three substituents selected from R^b , OH, (C_1-C_6) alkoxy, halogen, CO_2H , CN, $O(C=O)C_1-C_6$ alkyl, oxo, NO_2 and $N(R^b)_2$;

30 R^8 and R^9 are independently selected from:

- 1) H,
- 2) $(C=O)O_bC_1-C_{10}$ alkyl,
- 3) $(C=O)O_bC_3-C_8$ cycloalkyl,
- 4) $(C=O)O_b$ aryl,
- 35 5) $(C=O)O_b$ heterocyclyl,

- 6) C₁-C₁₀ alkyl,
- 7) aryl,
- 8) C₂-C₁₀ alkenyl,
- 9) C₂-C₁₀ alkynyl,
- 10) heterocyclyl,
- 11) C₃-C₈ cycloalkyl,
- 12) SO₂R^a, and
- 13) (C=O)NR^b₂,

said alkyl, cycloalkyl, aryl, heterocyclyl, alkenyl, and alkynyl is optionally substituted with one, two or three substituents selected from R⁷, or

R⁸ and R⁹ can be taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 3-7 members in each ring and optionally containing, in addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with one, two or three substituents selected from R⁷;

R¹² is selected from: F and -CH₂F;

R¹⁴ is selected from: H and -CH₂F, provided that when t is 1, R¹⁴ is H;

R^a is independently selected from: (C₁-C₆)alkyl, (C₃-C₆)cycloalkyl, aryl, or heterocyclyl, optionally substituted with one, two or three substituents selected from R⁷;

R^b is independently selected from: H, (C₁-C₆)alkyl, aryl, heterocyclyl, (C₃-C₆)cycloalkyl, (C=O)OC₁-C₆ alkyl, (C=O)C₁-C₆ alkyl, (C=O)aryl, (C=O)heterocyclyl, (C=O)NR^eRe' or S(O)₂R^a, optionally substituted with one, two or three substituents selected from R⁷;

R^c and R^c' are independently selected from: H, (C₁-C₆)alkyl, aryl, NH₂, OH, OR^a, -(C₁-C₆)alkyl-OH, -(C₁-C₆)alkyl-O-(C₁-C₆)alkyl, (C=O)OC₁-C₆ alkyl, (C=O)C₁-C₆ alkyl, (C=O)aryl, (C=O)heterocyclyl, (C=O)NR^eRe', S(O)₂R^a and -(C₁-C₆)alkyl-N(R^b)₂, wherein the alkyl is optionally substituted with one, two or three substituents selected from R⁷; or


R^c and R^c' can be taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 3-7 members in each ring and optionally containing, in

addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with one, two or three substituents selected from R⁷;

- 5 R^e and R^{e'} are independently selected from: H, (C₁-C₆)alkyl, aryl, heterocyclyl and (C₃-C₆)cycloalkyl, optionally substituted with one, two or three substituents selected from R⁷; or

- R^e and R^{e'} can be taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 3-7 members in each ring and optionally containing, in
 10 addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with one, two or three substituents selected from R⁷.

4. The compound according to Claim 3 of the Formula IV:



IV

or a pharmaceutically acceptable salt or stereoisomer thereof,

wherein:

- 20 a is 0 or 1;
 b is 0 or 1;
 m is 0, 1, or 2;
 r is 0 or 1;

s is 0 or 1;

R¹ and R² are independently selected from: H and (C₁-C₆)alkyl, optionally substituted with one, two or three substituents selected from R⁷;

5

R⁴ is independently selected from:

- 1) halo,
- 2) OH,
- 3) O_bC₁-C₆ perfluoroalkyl,

10

R⁷ is selected from:

- 1) (C=O)_rO_s(C₁-C₁₀)alkyl,
- 2) O_r(C₁-C₃)perfluoroalkyl,
- 3) oxo,
- 15 4) OH,
- 5) halo,
- 6) CN,
- 7) (C₂-C₁₀)alkenyl,
- 8) (C₂-C₁₀)alkynyl,
- 20 9) (C=O)_rO_s(C₃-C₆)cycloalkyl,
- 10) (C=O)_rO_s(C₀-C₆)alkylene-aryl,
- 11) (C=O)_rO_s(C₀-C₆)alkylene-heterocyclyl,
- 12) (C=O)_rO_s(C₀-C₆)alkylene-N(R^b)₂,
- 13) C(O)R^a,
- 25 14) (C₀-C₆)alkylene-CO₂R^a,
- 15) C(O)H,
- 16) (C₀-C₆)alkylene-CO₂H, and
- 17) C(O)N(R^b)₂,
- 18) S(O)_mR^a, and
- 30 19) S(O)₂N(R^b)₂;

said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, alkylene and heterocyclyl is optionally substituted with up to three substituents selected from R^b, OH, (C₁-C₆)alkoxy, halogen, CO₂H, CN, O(C=O)C₁-C₆ alkyl, oxo, NO₂ and N(R^b)₂;

35 R⁸ and R⁹ are independently selected from:

- 1) H,
- 2) (C=O)O_bC₁-C₁₀ alkyl,
- 3) (C=O)O_bC₃-C₈ cycloalkyl,
- 4) (C=O)O_baryl,
- 5) (C=O)O_bheterocyclyl,
- 6) C₁-C₁₀ alkyl,
- 7) aryl,
- 8) C₂-C₁₀ alkenyl,
- 9) C₂-C₁₀ alkynyl,
- 10) heterocyclyl,
- 11) C₃-C₈ cycloalkyl,
- 12) SO₂R^a, and
- 13) (C=O)NR^b₂,

said alkyl, cycloalkyl, aryl, heterocyclyl, alkenyl, and alkynyl is optionally substituted with one, two or three substituents selected from R⁷, or

R⁸ and R⁹ can be taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 3-7 members in each ring and optionally containing, in addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with one, two or three substituents selected from R⁷;

R^a is independently selected from: (C₁-C₆)alkyl, (C₃-C₆)cycloalkyl, aryl, or heterocyclyl, optionally substituted with one, two or three substituents selected from R⁷;

R^b is independently selected from: H, (C₁-C₆)alkyl, aryl, heterocyclyl, (C₃-C₆)cycloalkyl, (C=O)OC₁-C₆ alkyl, (C=O)C₁-C₆ alkyl, (C=O)aryl, (C=O)heterocyclyl, (C=O)NR^eR^e ' or S(O)₂R^a, optionally substituted with one, two or three substituents selected from R⁷;

R^c and R^c ' are independently selected from: H, (C₁-C₆)alkyl, aryl, NH₂, OH, OR^a, -(C₁-C₆)alkyl-OH, -(C₁-C₆)alkyl-O-(C₁-C₆)alkyl, (C=O)OC₁-C₆ alkyl, (C=O)C₁-C₆ alkyl, (C=O)aryl, (C=O)heterocyclyl, (C=O)NR^eR^e ', S(O)₂R^a and -(C₁-C₆)alkyl-N(R^b)₂, wherein the alkyl is optionally substituted with one, two or three substituents selected from R⁷; or

R^c and $R^{c'}$ can be taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 3-7 members in each ring and optionally containing, in addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with one, two or three substituents selected from R^7 ;

R^e and $R^{e'}$ are independently selected from: H, (C_1-C_6) alkyl, aryl, heterocyclyl and (C_3-C_6) cycloalkyl, optionally substituted with one, two or three substituents selected from R^7 ; or

R^e and $R^{e'}$ can be taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 3-7 members in each ring and optionally containing, in addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with one, two or three substituents selected from R^7 .


5. A compound selected from:

(2*S*)-4-(2,5-difluorophenyl)-*N*-[(4*R*,6*S*)-6-fluoro-1-methylazepan-4-yl]-2-(hydroxymethyl)-*N*-methyl-2-phenyl-2,5-dihydro-1*H*-pyrrole-1-carboxamide








(2*S*)-4-(2,5-difluorophenyl)-*N*-[(4*S*,6*R*)-6-fluoro-1-methylazepan-4-yl]-2-(hydroxymethyl)-*N*-methyl-2-phenyl-2,5-dihydro-1*H*-pyrrole-1-carboxamide








or a pharmaceutically acceptable salt thereof.











6. The compound according to Claim 1 which is selected from:















R ₁	R ₂	R ₃	R ₄	R ₅
	CH ₂ OH	Me	F	H
	CH ₂ OH	Me	F	H
	CH ₂ OH	Me	F	H
	CH ₂ OH	Me	F	H
	CH ₂ OH	Me	F	H
	CH ₂ OH	Me	F	H
	CH ₂ OH	Me	F	H









R ₁	R ₂	R ₃	R ₄	R ₅
	CH ₂ OH	Me	F	H
	CH ₂ OH	Me	F	H
	CH ₂ OH	Me	F	H
	CH ₂ OH	Me	F	H
	CH ₂ OH	Me	F	H
	CH ₂ OH	Me	F	H
	CH ₂ OH	Me	F	H

R ₁	R ₂	R ₃	R ₄	R ₅
	CH ₂ OH	Me	F	H
	CH ₂ OH	Me	F	H
	CH ₂ OH	Me	F	H
	CH ₂ OH	Me	F	H
	CH ₂ OH	Me	F	H
Me	Me	Me	F	H
Me		Me	F	H
Me		Me	F	H


R ₁	R ₂	R ₃	R ₄	R ₅
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H

R ₁	R ₂	R ₃	R ₄	R ₅
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H








R ₁	R ₂	R ₃	R ₄	R ₅
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me	CH ₂ OH		F	H
Me	CH ₂ OH		F	H








R ₁	R ₂	R ₃	R ₄	R ₅
Me	CH ₂ OH		F	H
Me	CH ₂ OH		F	H
Me	CH ₂ OH		F	H
Me	CH ₂ OH		F	H
Me	CH ₂ OH		F	H
Me	CH ₂ OH		F	H
Me	CH ₂ OH		F	H
Me	CH ₂ OH		F	H











R ₁	R ₂	R ₃	R ₄	R ₅
Me	CH ₂ OH	Me	Cl	H
Me	CH ₂ OH	Me	Br	H
Me	CH ₂ OH	Me	CN	H
Me	CH ₂ OH	Me	Me	H
Me	CH ₂ OH	Me	CF ₃	H
Me	CH ₂ OH	Me	NO ₂	H
Me	CH ₂ OH	Me	F	OH
Me	CH ₂ OH	Me	F	NH ₂
Me	CH ₂ OH	Me	F	F
Me	CH ₂ OH	Me	F	SH















R ₁	R ₂	R ₃	R ₄	R ₅
	CH ₂ OH	Me	F	H
	CH ₂ OH	Me	F	H
	CH ₂ OH	Me	F	H
	CH ₂ OH	Me	F	H
	CH ₂ OH	Me	F	H
	CH ₂ OH	Me	F	H
	CH ₂ OH	Me	F	H









R ₁	R ₂	R ₃	R ₄	R ₅
	CH ₂ OH	Me	F	H
	CH ₂ OH	Me	F	H
	CH ₂ OH	Me	F	H
	CH ₂ OH	Me	F	H
	CH ₂ OH	Me	F	H
	CH ₂ OH	Me	F	H
	CH ₂ OH	Me	F	H

R ₁	R ₂	R ₃	R ₄	R ₅
	CH ₂ OH	Me	F	H
	CH ₂ OH	Me	F	H
	CH ₂ OH	Me	F	H
	CH ₂ OH	Me	F	H
	CH ₂ OH	Me	F	H
Me	Me	Me	F	H
Me		Me	F	H
Me		Me	F	H


R ₁	R ₂	R ₃	R ₄	R ₅
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H

R ₁	R ₂	R ₃	R ₄	R ₅
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H








R ₁	R ₂	R ₃	R ₄	R ₅
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me	CH ₂ OH		F	H
Me	CH ₂ OH		F	H








R ₁	R ₂	R ₃	R ₄	R ₅
Me	CH ₂ OH		F	H
Me	CH ₂ OH		F	H
Me	CH ₂ OH		F	H
Me	CH ₂ OH		F	H
Me	CH ₂ OH		F	H
Me	CH ₂ OH		F	H
Me	CH ₂ OH		F	H
Me	CH ₂ OH		F	H











R ₁	R ₂	R ₃	R ₄	R ₅
Me	CH ₂ OH	Me	Cl	H
Me	CH ₂ OH	Me	Br	H
Me	CH ₂ OH	Me	CN	H
Me	CH ₂ OH	Me	Me	H
Me	CH ₂ OH	Me	CF ₃	H
Me	CH ₂ OH	Me	NO ₂	H
Me	CH ₂ OH	Me	F	OH
Me	CH ₂ OH	Me	F	NH ₂
Me	CH ₂ OH	Me	F	F
Me	CH ₂ OH	Me	F	SH















R ₁	R ₂	R ₃	R ₄	R ₅
	CH ₂ OH	Me	F	H
	CH ₂ OH	Me	F	H
	CH ₂ OH	Me	F	H
	CH ₂ OH	Me	F	H
	CH ₂ OH	Me	F	H
	CH ₂ OH	Me	F	H
	CH ₂ OH	Me	F	H









R ₁	R ₂	R ₃	R ₄	R ₅
	CH ₂ OH	Me	F	H
	CH ₂ OH	Me	F	H
	CH ₂ OH	Me	F	H
	CH ₂ OH	Me	F	H
	CH ₂ OH	Me	F	H
	CH ₂ OH	Me	F	H
	CH ₂ OH	Me	F	H

R ₁	R ₂	R ₃	R ₄	R ₅
	CH ₂ OH	Me	F	H
	CH ₂ OH	Me	F	H
	CH ₂ OH	Me	F	H
	CH ₂ OH	Me	F	H
	CH ₂ OH	Me	F	H
Me	Me	Me	F	H
Me		Me	F	H
Me		Me	F	H


R ₁	R ₂	R ₃	R ₄	R ₅
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H








R ₁	R ₂	R ₃	R ₄	R ₅
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H








R ₁	R ₂	R ₃	R ₄	R ₅
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me	CH ₂ OH		F	H
Me	CH ₂ OH		F	H








R_1	R_2	R_3	R_4	R_5
Me	CH ₂ OH		F	H
Me	CH ₂ OH		F	H
Me	CH ₂ OH		F	H
Me	CH ₂ OH		F	H
Me	CH ₂ OH		F	H
Me	CH ₂ OH		F	H
Me	CH ₂ OH		F	H
Me	CH ₂ OH		F	H











R ₁	R ₂	R ₃	R ₄	R ₅
Me	CH ₂ OH	Me	Cl	H
Me	CH ₂ OH	Me	Br	H
Me	CH ₂ OH	Me	CN	H
Me	CH ₂ OH	Me	Me	H
Me	CH ₂ OH	Me	CF ₃	H
Me	CH ₂ OH	Me	NO ₂	H
Me	CH ₂ OH	Me	F	OH
Me	CH ₂ OH	Me	F	NH ₂
Me	CH ₂ OH	Me	F	F
Me	CH ₂ OH	Me	F	SH















R_1	R_2	R_3	R_4	R_5
	CH ₂ OH	Me	F	H
	CH ₂ OH	Me	F	H
	CH ₂ OH	Me	F	H
	CH ₂ OH	Me	F	H
	CH ₂ OH	Me	F	H
	CH ₂ OH	Me	F	H
	CH ₂ OH	Me	F	H









R ₁	R ₂	R ₃	R ₄	R ₅
	CH ₂ OH	Me	F	H
	CH ₂ OH	Me	F	H
	CH ₂ OH	Me	F	H
	CH ₂ OH	Me	F	H
	CH ₂ OH	Me	F	H
	CH ₂ OH	Me	F	H
	CH ₂ OH	Me	F	H

R ₁	R ₂	R ₃	R ₄	R ₅
	CH ₂ OH	Me	F	H
	CH ₂ OH	Me	F	H
	CH ₂ OH	Me	F	H
	CH ₂ OH	Me	F	H
	CH ₂ OH	Me	F	H
Me	Me	Me	F	H
Me		Me	F	H
Me		Me	F	H

R ₁	R ₂	R ₃	R ₄	R ₅
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H

R ₁	R ₂	R ₃	R ₄	R ₅
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H

R ₁	R ₂	R ₃	R ₄	R ₅
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me		Me	F	H
Me	CH ₂ OH		F	H
Me	CH ₂ OH		F	H

R ₁	R ₂	R ₃	R ₄	R ₅
Me	CH ₂ OH		F	H
Me	CH ₂ OH		F	H
Me	CH ₂ OH		F	H
Me	CH ₂ OH		F	H
Me	CH ₂ OH		F	H
Me	CH ₂ OH		F	H
Me	CH ₂ OH		F	H
Me	CH ₂ OH		F	H

R ₁	R ₂	R ₃	R ₄	R ₅
Me	CH ₂ OH	Me	Cl	H
Me	CH ₂ OH	Me	Br	H
Me	CH ₂ OH	Me	CN	H
Me	CH ₂ OH	Me	Me	H
Me	CH ₂ OH	Me	CF ₃	H
Me	CH ₂ OH	Me	NO ₂	H
Me	CH ₂ OH	Me	F	OH
Me	CH ₂ OH	Me	F	NH ₂
Me	CH ₂ OH	Me	F	F
Me	CH ₂ OH	Me	F	SH

or a pharmaceutically acceptable salt or stereoisomer thereof.

7. A pharmaceutical composition that is comprised of a compound in accordance with Claim 1 and a pharmaceutically acceptable carrier.

5

8. A method of treating or preventing cancer in a mammal in need of such treatment that is comprised of administering to said mammal a therapeutically effective amount of a compound of Claim 1.

10 9. A method of treating cancer or preventing cancer in accordance with Claim 8 wherein the cancer is selected from cancers of the brain, genitourinary tract, lymphatic system, stomach, larynx and lung.

15 10. A method of treating or preventing cancer in accordance with Claim 8 wherein the cancer is selected from histiocytic lymphoma, lung adenocarcinoma, small cell lung cancers, pancreatic cancer, glioblastomas and breast carcinoma.

20 11. A process for making a pharmaceutical composition which comprises combining a compound of Claim 1 with a pharmaceutically acceptable carrier.

12. The composition of Claim 7 further comprising a second compound selected from:

- 25
- 1) an estrogen receptor modulator,
 - 2) an androgen receptor modulator,
 - 3) a retinoid receptor modulator,
 - 4) a cytotoxic/cytostatic agent,
 - 5) an antiproliferative agent,
 - 6) a prenyl-protein transferase inhibitor,
 - 7) an HMG-CoA reductase inhibitor,
 - 30 8) an HIV protease inhibitor,
 - 9) a reverse transcriptase inhibitor,
 - 10) an angiogenesis inhibitor,
 - 11) a PPAR- γ agonist,
 - 12) a PPAR- δ agonists;
 - 35 13) an inhibitor of cell proliferation and survival signaling,

- 14) an agent that interferes with a cell cycle checkpoint, and
- 15) an apoptosis inducing agent.

13. The composition of Claim 12, wherein the second compound is an
5 angiogenesis inhibitor selected from the group consisting of a tyrosine kinase inhibitor, an
inhibitor of epidermal-derived growth factor, an inhibitor of fibroblast-derived growth factor, an
inhibitor of platelet derived growth factor, an MMP inhibitor, an integrin blocker, interferon- α ,
interleukin-12, pentosan polysulfate, a cyclooxygenase inhibitor, carboxyamidotriazole,
combretastatin A-4, squalamine, 6-O-(chloroacetyl-carbonyl)-fumagillol, thalidomide,
10 angiostatin, troponin-1, and an antibody to VEGF.

14. The composition according to Claim 7 further comprising a proteasome
inhibitor.

15. The composition according to Claim 7 further comprising a aurora kinase
inhibitor.

16. The composition according to Claim 7 further comprising a Raf kinase
inhibitor.

17. The composition according to Claim 7 further comprising a
serine/threonine kinase inhibitor.

18. The composition according to Claim 7 further comprising an inhibitor of
25 another mitotic kinesin which is not KSP.

19. The composition of Claim 12, wherein the second compound is an
estrogen receptor modulator selected from tamoxifen and raloxifene.

20. A method of treating cancer which comprises administering a
30 therapeutically effective amount of a compound of Claim 1 in combination with radiation
therapy.

21. A method of treating or preventing cancer that comprises administering a therapeutically effective amount of a compound of Claim 1 in combination with a compound selected from:

- 1) an estrogen receptor modulator,
- 2) an androgen receptor modulator,
- 3) a retinoid receptor modulator,
- 4) a cytotoxic/cytostatic agent,
- 5) an antiproliferative agent,
- 6) a prenyl-protein transferase inhibitor,
- 7) an HMG-CoA reductase inhibitor,
- 8) an HIV protease inhibitor,
- 9) a reverse transcriptase inhibitor,
- 10) an angiogenesis inhibitor,
- 11) PPAR- γ agonists,
- 12) PPAR- δ agonists,
- 13) an inhibitor of inherent multidrug resistance,
- 14) an anti-emetic agent,
- 15) an agent useful in the treatment of anemia,
- 16) an agent useful in the treatment of neutropenia,
- 17) an immunologic-enhancing drug,
- 18) an inhibitor of cell proliferation and survival signaling,
- 19) an agent that interferes with a cell cycle checkpoint, and
- 20) an apoptosis inducing agent.

22. A method of treating cancer that comprises administering a therapeutically effective amount of a compound of Claim 1 in combination with radiation therapy and a compound selected from:

- 1) an estrogen receptor modulator,
- 2) an androgen receptor modulator,
- 3) a retinoid receptor modulator,
- 4) a cytotoxic/cytostatic agent,
- 5) an antiproliferative agent,
- 6) a prenyl-protein transferase inhibitor,
- 7) an HMG-CoA reductase inhibitor,
- 8) an HIV protease inhibitor,

- 5 9) a reverse transcriptase inhibitor,
 10) an angiogenesis inhibitor,
 11) PPAR- γ agonists,
 12) PPAR- δ agonists,
 13) an inhibitor of inherent multidrug resistance,
 14) an anti-emetic agent,
 15) an agent useful in the treatment of anemia,
 16) an agent useful in the treatment of neutropenia,
 17) an immunologic-enhancing drug,
10 18) an inhibitor of cell proliferation and survival signaling,
 19) an agent that interferes with a cell cycle checkpoint, and
 20) an apoptosis inducing agent.
- 15 23. A method of treating or preventing cancer which comprises administering
a therapeutically effective amount of a compound of Claim 1 and paclitaxel or trastuzumab.
24. A method of treating or preventing cancer which comprises administering
a therapeutically effective amount of a compound of Claim 1 and a GPIIb/IIIa antagonist.
- 20 25. The method of Claim 24 wherein the GPIIb/IIIa antagonist is tirofiban.
26. A method of treating or preventing cancer which comprises administering
a therapeutically effective amount of a compound of Claim 1 in combination with a COX-2
inhibitor.
- 25 27. A method of treating or preventing cancer which comprises administering
a therapeutically effective amount of a compound of Claim 1 in combination with a proteasome
inhibitor.
- 30 28. A method of treating or preventing cancer which comprises administering
a therapeutically effective amount of a compound of Claim 1 in combination with an aurora
kinase inhibitor.

29. A method of treating or preventing cancer which comprises administering a therapeutically effective amount of a compound of Claim 1 in combination with a Raf kinase inhibitor.

5 30. A method of treating or preventing cancer which comprises administering a therapeutically effective amount of a compound of Claim 1 in combination with a serine/threonine kinase inhibitor.

10 31. A method of treating or preventing cancer which comprises administering a therapeutically effective amount of a compound of Claim 1 in combination with an inhibitor of a mitotic kinesin that is not KSP.

15 32. A method of modulating mitotic spindle formation which comprises administering a therapeutically effective amount of a compound of Claim 1.

33. A method of inhibiting the mitotic kinesin KSP which comprises administering a therapeutically effective amount of a compound of Claim 1.